

DOCTORS FOSTER^{and}SMITH[®] PHARMACY

Patient Information Sheet

PO Box 100, Rhinelander WI 54501
For questions or to consult with a pharmacist, call 1-800-447-3021 Mon-Fri from 9am-6pm CST and Sat from 9am-1pm CST. If you have an emergency situation, please contact your local veterinarian immediately.

Imidacloprid + Moxidectin

(Advantage Multi)

Common Drug Name

Imidacloprid plus moxidectin

Common Brand Names

Advantage Multi for Cats

Advantage Multi for Dogs

Generic products are not available.

Storage

Store at room temperature away from children and pets.

Uses

The combination of imidacloprid and moxidectin is used in dogs and cats for the prevention of heartworm disease and the treatment and control of flea infestations, hookworms and roundworms. It is also used for the treatment and control of ear mites in cats and whipworms in dogs.

Dose and Administration

Always follow the dosage instructions provided by your veterinarian. If you have difficulty giving the medication, contact your veterinarian.

The combination of imidacloprid and moxidectin is applied topically (on the skin). It is NOT given orally. Follow the manufacturer's recommendations for application. Do not apply to irritated skin.

If you miss a dose, give it as soon as you remember. If it is almost time for the next dose, skip the one you missed and go back to the regular schedule. Do not give 2 doses at once.

This medication should only be given to the pet for whom it was prescribed.

Possible Side Effects

Potential side effects in **dogs** include scratching, rubbing, or licking the application site; lethargy, decreased appetite, or hyperactivity. Consult your veterinarian if you notice any of the above

side effects. A medicinal odor and residue at the application site may also be seen. If a dog ingests the product or licks the application site, serious adverse effects may occur including depression, salivation, incoordination, panting and tremors.

Potential side effects in **cats** include lethargy and prolonged periods of sleep; behavioral changes including agitation, excessive grooming, hiding, pacing; etc.; discomfort at the application site (rubbing, scratching, etc.); increased salivation within 1 hour of treatment; increased drinking; or coughing. If cat licks the application site may see increased salivation, tremors, vomiting, and decreased appetite.

Consult your veterinarian if you notice any of the above side effects in your pet.

Hair may appear stiff, matted, oily, or damp after application.

To report a possible adverse reaction to this product, contact the manufacturer (Bayer Health Care) at 1-800-422-9874. For questions, call the manufacturer at 1-800-255-6826.

If your pet experiences an allergic reaction to the medication, signs may include facial swelling, hives, scratching, sudden onset of diarrhea, vomiting, shock, seizures, pale gums, cold limbs, or coma. If you observe any of these signs, contact your veterinarian immediately.

Precautions

Precautions in Animals:

Not for use in animals hypersensitive (allergic) to imidacloprid or moxidectin.

Do not use in animals who are sick, debilitated, or underweight.

Do not use in cats under 9 weeks of age or less than 2 pounds, or in dogs under 7 weeks of age or less than 3 pounds.

The safety of imidacloprid plus moxidectin in breeding, pregnant or lactating animals (female animals nursing their young) has not been determined.

The safety of imidacloprid plus moxidectin in geriatric animals with potential health problems has not been determined.

Do not use the dog product on cats.

Do not allow any animal to lick the application site for 30 minutes. Separation of treated animals may be necessary.

Consult with your veterinarian regarding the physical examinations and laboratory testing necessary prior to and during treatment with imidacloprid plus moxidectin.

Human Precautions:

Children should not come in contact with the application site for 30 minutes after application.

Can cause eye irritation and be harmful if swallowed. Wash hands thoroughly with soap and water after handling.

Drug, Food and Test Interactions

Notify your veterinarian of any other medications, including vitamins, supplements, and other flea/tick products your pet is receiving while using imidacloprid and moxidectin on your pet.

Signs of Toxicity/Overdose

May include depression, increased salivation, tremors, or incoordination. If you know or suspect your pet has had an overdose, or if you observe any of these signs in your pet, contact your veterinarian immediately.

Keep this and all other medications out of the reach of children and pets.